

# Recombinant Mouse KIRREL1/NEPH1 Protein (His Tag)



Catalog Number:PKSM040812

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Kin of IRRE-like protein 1;Kin of irregular chiasm-like protein 1;Nephrin-like protein 1;Kirrel1;Neph1
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Leu 525
<b>Accession</b>	NP_570937.2
<b>Calculated Molecular Weight</b>	53.8 kDa
<b>Observed molecular weight</b>	65-70 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Measured by the ability of the immobilized protein to support the adhesion of MS1 mouse pancreatic islet endothelial cells (ATCC: CRL2279). When cells are added to coated plates (30 µg/mL, 100 µL/well), > 40% cells will adhere specifically after 90 minutes at 37 °C.

## Properties

<b>Purity</b>	> 98 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 98 % as determined by reducing SDS-PAGE.

## Background

NEPH1 (KIRREL1) belongs to a family of three closely related transmembrane proteins of the Ig superfamily with a structure similar to that of nephrin. All three Neph proteins share a conserved podocin-binding motif; mutation of a centrally located tyrosine residue dramatically lowers the affinity of Neph1 for podocin. Neph1 triggers AP-1 activation

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse KIRREL1/NEPH1 Protein (His Tag)



Catalog Number:PKSM040812

similarly to nephrin but requires the presence of Tec family kinases for efficient transactivation. Nep1 consists of a signal peptide, five Ig-like C2-type domains with the middle domain overlapping with a PKD-like domain, an RGD sequence, a transmembrane domain and a cytoplasmic tail, which is expressed in slit diaphragm domains of podocytes and in vertebrate and invertebrate nervous systems. Nep1 is abundantly expressed in the kidney, specifically expressed in podocytes of kidney glomeruli, and plays a significant role in the normal development and function of the glomerular permeability. Nep1 interacts with nephrin in vitro and in vivo, and able to stimulate transcriptional activation in a model system, such as the activation the transcription factor AP-1 via the stimulation of a MAPK module. Nep1 is crucial for the integrity of the slit diaphragm, as Nep1 gene knockout mice results in effacement of glomerular podocytes, heavy proteinuria, and early postnatal death.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)