Recombinant SARS-CoV Spike Protein (RBD, mFc Tag)

Catalog Number: PKSV030106



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms coronavirus s1 Protein; SARS; coronavirus s2 Protein; SARS; coronavirus spike

Protein;SARS;cov spike Protein;SARS;ncov RBD Protein;SARS;ncov s1 Protein;SARS;ncov s2 Protein;SARS;ncov spike Protein;SARS;novel coronavirus

RBD Protein;SARS;novel coronavirus s1 Protein;SARS;novel coronavirus s2 Protein;SARS;novel coronavirus spike Protein;SARS;novel coronavirus s2

Protein;SARS;s2 Protein;SARS;Spike RBD Protein;SARS

Species SARS

Expression Host HEK293 Cells
Sequence Arg306-Phe527
Accession AAX16192.1
Calculated Molecular Weight 51.4 kDa
Tag C-mFc1

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as


protectants before lyophilization.


Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data







Immobilized human ACE2 protein (His tag) at 2μg/mL (100μL/well) can bind SARS-CoV Spike/RBD Protein (RBD, mFc Tag), the EC50 of SARS-CoV Spike/RBD Protein (RBD, mFc Tag) is 9-40 ng/mL.

Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com
Email: techsupport@elabscience.com

Recombinant SARS-CoV Spike Protein (RBD, mFc Tag)

Catalog Number: PKSV030106



Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com