# Recombinant Mouse TNFRSF1B/CD120b Protein (Fc Tag)

**Catalog No.** PKSM040847

***Note:*** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

| | |
|---|---|
| **Synonyms** | CD120b;p75;TNF-alphaR2;TNF-R-II;TNF-R2;TNF-R75;TNFalpha-R2;TNFBR;Tnfr-1;Tnfr2;TNFR80;TNFRII;Tumor necrosis factor receptor superfamily member 1b;Tnfrsf1b |
| **Species** | Mouse |
| **Expression Host** | HEK293 Cells |
| **Sequence** | Met 1-Gly 258 |
| **Accession** | NP_035740.2 |
| **Calculated Molecular Weight** | 52.3 kDa |
| **Observed molecular weight** | 65 kDa |
| **Tag** | C-hFc |
| **Bioactivity** | Measured in a cytotoxicity assay using L929 mouse fibrosarcoma cells in the presence of the metabolic inhibitor actinomycin D. The ED50 for this effect is typically 20-80 ng/mL in the presence of 0.5 ng/mL of Recombinant Mouse TNFα. |

## Properties

| | |
|---|---|
| **Purity** | > 95 % as determined by reducing SDS-PAGE. |
| **Endotoxin** | < 1.0 EU per μg of the protein as determined by the LAL method. |
| **Storage** | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months. |
| **Shipping** | This product is provided as lyophilized powder which is shipped with ice packs. |
| **Formulation** | Lyophilized from sterile PBS, pH 7.4<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| **Reconstitution** | Please refer to the printed manual for detailed information. |

## Data




> 95 % as determined by reducing SDS-PAGE.

## Background

Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B), also known as Tumor necrosis factor receptor 2 (TNFR2) or CD120b antigen, is a member of the tumor necrosis factor receptor superfamily. TNFR2/CD120b/TNFRSF1B is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. TNFR2/CD120b/TNFRSF1B is not a major contributing factor to the genetic risk of type 2 diabetes, its associated peripheral neuropathy and hypertension and related metabolic traits in North Indians. Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B) has been reported to be associated with SLE risk in Japanese populations. TNFR2/CD120b/TNFRSF1B serves as a receptor with high affinity for TNFSF2 and approximately 5-fold lower affinity for homotrimeric TNFSF1. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.


Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com