

Recombinant Mouse SIGIRR/TIR8 Protein (His & Fc Tag)



Catalog Number:PKSM040853

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI256711;TIR8
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-His 117
Accession	NP_075546.2
Calculated Molecular Weight	40.7 kDa
Observed molecular weight	60-65 kDa
Tag	C-His-Fc

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Single Ig IL-1-related receptor (SIGIRR) or TIR8 is a member of Toll-like receptor-interleukin 1 receptor signaling (TLR-IL-1R) receptor superfamily. Although SIGIRR/TIR8 shows the typical conserved motifs that characterize the IL-1R and Toll superfamily, it is structurally and functionally distinct from both. SIGIRR/TIR8 has only one Ig domain in its extracellular portion whereas the IL-1R family contains three Ig folds. An unusually long cytoplasmic domain is reminiscent of the structure of drosophila Toll, yet the SIGIRR peptide sequence is more closely related to IL-1RI. SIGIRR/TIR8 was mainly expressed in mouse and human epithelial tissues such as kidney, lung and gut. Resting and activated T and B lymphocytes and monocytes-macrophages expressed little or no SIGIRR/TIR8, with the exception of the mouse GG2EE macrophage line. Inflammation is enhanced in SIGIRR-deficient mice. SIGIRR negatively modulates

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse SIGIRR/TIR8 Protein (His & Fc Tag)



Catalog Number: PKSM040853

immune responses. Inflammation is enhanced in SIGIRR-deficient mice, as shown by their enhanced chemokine induction after IL-1 injection and reduced threshold for lethal endotoxin challenge.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com