

Recombinant Mouse Latexin/LXN Protein (His Tag)

Catalog No. PKSM040884

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	MGC144352;MGC144353
Species	Mouse
Expression Host	E.coli
Sequence	Glu 2-Glu 222
Accession	NP_058033.2
Calculated Molecular Weight	26.3 kDa
Observed molecular weight	32 kDa
Tag	N-His
Bioactivity	Measured by its ability to inhibit carboxypeptidase-A1 cleavage of the colorimetric peptide substrate Ac-Phe-Thiaphe-OH in the presence of 5, 5'Dithiobis(2-nitrobenzoic acid) (DTNB). The IC50 value is < 2.0 nM.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 8.0, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Latexin, also known as endogenous carboxypeptidase inhibitor, tissue carboxypeptidase inhibitor, TCI, ECI and LXN, is a cytoplasm protein which belongs to the protease inhibitor I47 (latexin) family. It is highly expressed in heart, prostate, ovary, kidney, pancreas, and colon. Latexin / LXN is the only known endogenous specific inhibitor of zinc-dependent metallopeptidases (MCPs) present in mammals so far. Latexin is originally identified as a molecular marker for the regional specification of the neocortex in development in rats. The 222 amino acid latexin in human shows different expression distribution with high levels in heart, prostate, ovary, kidney, pancreas, and colon, but only moderate or low levels in other tissues including brain. Latexin is also expressed at high levels and is inducible in macrophages in concert with other protease inhibitors and potential protease targets, and thus is suggested to play a role in inflammation and innate immunity pathways. Despite of the non-detectable sequence similarity with plant and parasite inhibitors, Latexin is related to a human putative tumor suppressor protein, TIG1. In addition, Latexin is also implicated in Alzheimer's disease.