

# Recombinant Mouse CD32/FCGR2B Protein (His & AVI Tag), Biotinylated

Catalog No. PKSM040900

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Synonyms AI528646;CD32;F630109E10Rik;Fcgr2;Fcgr2a;FcgRII;Fcr-2;Fcr-3;fcRII;Fc[g]RII;

Ly-17;Ly-m20;LyM-1

Species Mouse

Expression HostHEK293 CellsSequenceMet 1-Arg 217AccessionNP\_001070657.1

Calculated Molecular Weight23.9 kDaObserved molecular weight38-43 kDaTagC-His-Avi

**Bioactivity** 1. Measured by its ability to bind human IgG1 in a functional ELISA. 2. Labeling

ratio of biotin to protein: 0.5

# **Properties**

**Purity** > 97 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

## Background

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Receptors for Fc portion of IgG (Fcγ Rs) are members of the Ig superfamily, and are divided into three classes designated Fcy RI (CD64), Fcy RII (CD32), and Fcy RIII (CD16). CD32 protein is a low affinity receptor for IgG that binds only IgG immune complexes and is expressed on a diverse range of cells such as monocytes, macrophages, neutrophils, eosinophils, platelets, and B cells. Human CD32 class is encoded by three closely related genes, and designated Fcy RII A, B, and C which share 94-99% amino acid identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. CD32 is involved in a number of immune responses including antibodydependent cell-mediated cytotoxicity, clearance of immune complexes, release of inflammatory mediators, and regulation of antibody production.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com