# **CD38 Polyclonal Antibody**

Catalog Number: E-AB-66945



Note: Centrifuge before opening to ensure complete recovery of vial contents.

# **Description**

Reactivity Human, Mouse, Rat

**Immunogen** Recombinant fusion protein of human CD38 (NP\_001766.2).

Host Rabbit
Isotype IgG

**Purification** Affinity purification


**Conjugation** Unconjugated

**Formulation** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


**Applications** Recommended Dilution

**IF** 1:50-1:200

### Data



Immunofluorescence analysis of THP-1 cells using CD38 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse spleen using CD38 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

# Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

# **Background**

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com