

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Rat
Immunogen	Recombinant human Serine/threonine-protein kinase 4 protein
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.

Applications Recommended Dilution

WB	1:500-1:1000
IHC	1:50-1:100


Data



www.elabscience.com

Western Blot analysis of 293T and Rat liver using STK4 Polyclonal Antibody at dilution of 1:600

Observed Mw:56 kDa
Calculated Mw:56 kDa



www.elabscience.com

Immunohistochemistry of paraffin-embedded Rat lung using STK4 Polyclonal Antibody at dilution of 1:50

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene is a cytoplasmic kinase that is structurally similar to the yeast Ste20p kinase, which acts upstream of the stress-induced mitogen-activated protein kinase cascade. The encoded protein can phosphorylate myelin basic protein and undergoes autophosphorylation. A caspase-cleaved fragment of the encoded protein has been shown to be capable of phosphorylating histone H2B. The particular phosphorylation catalyzed by this protein has been correlated with apoptosis, and it's possible that this protein induces the chromatin condensation observed in this process.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017