

A Reliable Research Partner in Life Science and Medicine

CBX3 Monoclonal Antibody

Catalog No. E-AB-22145

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Recombinant Protein of HP-1γ

Host Mouse Isotype IgG

Purification Protein A purification


Conjugation Unconjugated

Buffer PBS with 0.02% sodium azide and 50% glycerol pH 7.4.

Applications Recommended Dilution


WB 1:500-2000 IHC 1:50-300

Data




Western Blot analysis of 1) Hela, 2)3T3, 3) PC-12 cells using CBX3 Monoclonal Antibody at dilution of 1:1000.

Observed Mw:24kDa



Immunohistochemistry of paraffin-embedded Human colon carcinoma tissue using CBX3 Monoclonal Antibody at dilution of 1:200.



Immunofluorescence analysis of Mouse spleen tissue using CBX3 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>





A Reliable Research Partner in Life Science and Medicine

Background

At the nuclear envelope, the nuclear lamina and heterochromatin are adjacent to the inner nuclear membrane. The protein encoded by this gene binds DNA and is a component of heterochromatin. This protein also can bind lamin B receptor, an integral membrane protein found in the inner nuclear membrane. The dual binding functions of the encoded protein may explain the association of heterochromatin with the inner nuclear membrane. This protein binds histone H3 tails methylated at Lys-9 sites. This protein is also recruited to sites of ultraviolet-induced DNA damage and double-strand breaks.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com