TPSD1 Polyclonal Antibody

Catalog No. E-AB-36155

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Synthesized peptide derived from Tryptase-3
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol pH 7.4.
Applications	Recommended Dilution
WB	1:500-2000
ELISA	1:10000-20000
Data	





Western Blot analysis of HepG2 cells using TPSD1 Polyclonal Antibody at dilution of 1:800. Observed Mw:26kDa Calculated Mw:27kDa Immunohistochemistry of paraffin-embedded Human tonsil tissue using TPSD1 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human skin tissue using TPSD1 Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

TPSD1 (Tryptase Delta 1) is a Protein Coding gene. GO annotations related to this gene include serine-type endopeptidase activity and serine-type peptidase activity. An important paralog of this gene is TPSAB1.Tryptases comprise a family of trypsin-like serine proteases, the peptidase family S1. Tryptases are enzymatically active only as heparin-stabilized tetramers, and they are resistant to all known endogenous proteinase inhibitors. Several tryptase genes are clustered on chromosome 16p13.3. These genes are characterized by several distinct features. They have a highly conserved 3' UTR and contain tandem repeat sequences at the 5' flank and 3' UTR which are thought to play a role in regulation of the mRNA stability. Although this gene may be an exception, most of the tryptase genes have an intron immediately upstream of the initiator Met codon, which separates the site of transcription initiation from protein coding sequence. This feature is characteristic of tryptases but is unusual in other genes. Tryptases have been implicated as mediators in the pathogenesis of asthma and other allergic and inflammatory disorders. This gene was once considered to be a pseudogene, although it is now believed to be a functional gene that encodes a protein.

For Research Use Only