

ZP2 Polyclonal Antibody

Catalog No. E-AB-63275

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Recombinant fusion protein of human ZP2 (NP_003451.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Applications Recommended Dilution

IF 1:50-1:200


Data




Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of Rat oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of Rat oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

For Research Use Only



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed of three glycoproteins with various functions during fertilization and preimplantation development. The glycosylated mature peptide is one of the structural components of the zona pellucida and functions in secondary binding and penetration of acrosome-reacted spermatozoa. Female mice lacking this gene do not form a stable zona matrix and are sterile. Alternative splicing results in multiple transcript variants.