ZP2 Polyclonal Antibody

Catalog Number: E-AB-63275



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Recombinant fusion protein of human ZP2 (NP_003451.1).

Host Rabbit
Isotype IgG

Purification Affinity purification


Conjugation Unconjugated

Formulation PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Applications Recommended Dilution

IF 1:50-1:200


Data



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of Rat oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100.

Blue: DAPI for nuclear staining.




Immunofluorescence analysis of Rat oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100.

Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse oophoroma cells using ZP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com
Email: techsupport@elabscience.com

ZP2 Polyclonal Antibody

Catalog Number: E-AB-63275



Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed of three glycoproteins with various functions during fertilization and preimplantation development. The glycosylated mature peptide is one of the structural components of the zona pellucida and functions in secondary binding and penetration of acrosome-reacted spermatozoa. Female mice lacking this gene do not form a stable zona matrix and are sterile. Alternative splicing results in multiple transcript variants.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>