Elabscience®

Elab Fluor[®] Violet 450 Anti-Mouse Ly6C Antibody[Monts 1]

Catalog No.E-AB-F1121UQStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityMouseApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

| Alternate Names | Lymphocyte antigen 6 complex, locus C,Ly 6C,Ly6c1,Ly6c2 |
|-----------------|--|
| Uniprot ID | P0CW03 |
| Gene ID | 17067 |
| Background | Most hematopoietic cells express one or more members of Ly-6 family. The expression of Ly-6 varies with development stage and activation. Ly-6C is a 14-17 kD GPI-linked surface protein |
| | expressed on mouse monocyte/macrophage cells, endothelial cells, neutrophils, and some T cell |
| | subsets. Ly-6C is reported to be an indicator of memory CD8+ T cells. |

Product Details

| Form | Liquid |
|---------------------|---|
| Concentration | 0.5 mg/mL |
| Size | 25µg/100µg |
| Clone No. | Monts 1 |
| Host | Rat |
| Isotype | Rat IgG2a, ĸ |
| Reactivity | Mouse |
| Application | FCM |
| Isotype Control | Elab Fluor [®] Violet 450 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833Q] |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |
| Shipping | Biological ice pack at 4 °C |
| Stability & Storage | Keep as concentrated solution. |
| | Store at 2~8°C and protected from prolonged exposure to light.Do not freeze. |
| | This product is guaranteed up to one year from purchase. |

For Research Use Only

Elabscience®

Fluorophore


Conjugation: Elab Fluor[®] Violet 450

Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is $0.1-1 \mu g/10^6$ cells in $100 \mu L$ volume].

Product data



C57BL/6 murine bone marrow cells are stained with Elab Fluor[®] Violet 450 Anti-Mouse Ly6C Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Related Information

- 1. Sample Preparation for Flow Cytometry <u>https://www.elabscience.com/List-detail-5594.html</u>
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>