A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD52 (C-Fc-Avi) Biotinylated

Catalog No. PKSH033863

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CAMPATH-1 antigen; CAMPATH-1; CD52 antigen; CD52

molecule;CD52;CDw52;Epididymal secretory protein E5;He5;HEL-

S-171mP;Human epididymis-specific protein 5

Species Human

Expression Host
Sequence
Gly25-Ser36
Accession
P31358
Calculated Molecular Weight
Observed molecular weight
Tag
HEK293 Cells
Gly25-Ser36
P31358
30.2 kDa
38-50 kDa
C-Fc-Avi

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

CD52, also known as CAMPATH-1 antigen, HE5, and gp20, is a cell surface glycoprotein that can be targeted to induce immune suppression by complement-mediated cell lysis. CD52 is expressed on lymphocytes, monocytes, monocytes derived dendritic cells, eosinophils, and neutrophils, as well as on mature spermatozoa and epithelial cells lining the male genital tract. From the clinical point of view this protein is an important target for therapeutic interventions aimed at leukocyte depletion in hematological malignancies and post-transplant immunosuppression. CD52 / CDW52 may play a role in carrying and orienting carbohydrate. It is an unusually good target for complement-mediated cell lysis.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com