A Reliable Research Partner in Life Science and Medicine

Recombinant Human Vitronectin/VTN (N-Truncated, C-6His)

Catalog No. PKSH033866

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Complement S-protein; epibolin; Serum Spreading Factor; Serum-spreading

factor; Somatomedin B; S-protein; V75; Vitronectin; VN; VNT; VTN

Species Human

Expression Host

Sequence

Val62-Leu478

Accession

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Val62-Leu478

AAH05046.1

48.3 kDa

60-70 kDa

C-His

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20 mMTris-HCl\25% Mannitol\250

mM NaCl20.02% Tween 802pH8.0


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Vitronectin, also known as VTN, is a large glycoprotein found in blood and the extracellular matrix (ECM). Vitronectin is a plasma glycoprotein implicated as a regulator of diverse physiological process, including blood coagulation, fibrinolysis, pericellular proteolysis, complement dependent immune responses, and cell attachment and spreading. Blocking of Hic(a member of the pneumococcal surface protein C (PspC) family) by specific antiserum or genetic deletion significantly reduced pneumococcal binding to soluble and immobilised vitronectin and to Factor H, respectively. In addition, Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com