

Recombinant Human Stanniocalcin 1/STC-1 (C-6His)

Catalog No. PKSH033893

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Stanniocalcin 1;stanniocalcin-1;STC1;STC-1;STCSTC-1
Species	Human
Expression Host	HEK293 Cells
Sequence	Thr18-Ala247
Accession	P52823
Calculated Molecular Weight	26.9 kDa
Observed molecular weight	28-36 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Stanniocalcin 1 (STC-1) is a homodimeric glycoprotein hormone that is involved in calcium and phosphate homeostasis. It was originally identified in bony fishes, where elevation of calcium in serum causes the release of STC from the

For Research Use Only

endocrine glands called the corpuscles of Stannius. STC-1 inhibits the breakdown of PAPP-A, protects cancer cells from apoptosis, reduces tumor size of liver cancers, promotes osteoblast differentiation and inhibits longitudinal bone growth directly at the growth plate. It is also a biomarker of brain and lung cancer progression. STC1 signals through inhibitory G-protein modulates CGRP receptor spatial localization during osteoblastogenesis and may function as a regulatory factor interacting with calcitonin peptide members during bone formation.

For Research Use Only