Recombinant Human FcRn & B2M Heterodimer (C-6His-Avi) Biotinylated



Catalog Number: PKSH033906

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	IgG receptor FcRn;Neonatal Fc receptor;FCRN	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Ala24-Ser297&Ile21-Met119	
Accession	AAF72596&P61769	
Calculated Molecular Weight	31.1&13.5 kDa	
Observed molecular weight	35&13 kDa	
Tag	C-His-Avi	
Bioactivity	Loaded Biotinylated Human FcRn Heterodimer-His-Avi on HIS1K Biosensor, can bind Anti-Human HER2 mAb with an affinity constant of 0.11 uM as determined in BLI assay.	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 6% Sucrose, 4% Mannitol, 0.05% Tween 80, pH 7.8. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer i	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

kDa	MK	R
120 90 60		
40	-	-
30	-	-
20	-	
14	-	

> 95 % as determined by reducing SDS-PAGE.

Background

FcRn complex consist of two subunits: IgG receptor FcRn large subunit p51^[2]alpha chain) and Beta-2-microglobulin^[2]Beta chain). The complexes is similar in structure to MHC class I-like heterodimer. Beta-2-microglobulin involved in the presentation of peptide antigens to the immune system. FcRn binds to the Fc region of monomeric immunoglobulins gamma, mediates the uptake of IgG from milk,Possible role in transfer of immunoglobulin G from mother to fetus. A

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Recombinant Human FcRn & B2M Heterodimer (C-6His-Avi) Biotinylated



Catalog Number: PKSH033906

principal component of antibody transport is the neonatal receptor for the Fc portion of immunoglobulin, a heterodimer of a MHC-1 alpha-chain homolog (FcRn) and beta-2-microglobulin (B2M).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com