Recombinant Human Leukocyte Ig-Like Receptor B2/LILRB2/ILT4/CD85d (C-Avi-6His) Biotinylated



Catalog Number: PKSH033919

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Leukocyte Immunoglobulin-Like Receptor Subfamily B Member

2;LIR-2;Leukocyte Immunoglobulin-Like Receptor 2;CD85 Antigen-Like Family Member D;Immunoglobulin-Like Transcript 4;ILT-4;Monocyte/Macrophage Immunoglobulin-Like Receptor 10;MIR-10;CD85d;LILRB2;ILT4;LIR2;MIR10

Species Human

Expression Host HEK293 Cells
Sequence Gln22-His458
Accession AAH36827.1
Calculated Molecular Weight 50.2 kDa
Observed molecular weight 60-80 kDa
Tag C-Avi-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Members of the immunoglobulin-like transcript (ILT) family are activating and inhibitory immunoreceptors whose genes are located same locus that encodes killer cell Ig-like receptors (KIR). Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2 (LIR-2) is a type I transmembrane protein. LIR-2 is expressed primarily on monocytes and dendritic cells (DC). Human LIR-2 is produced as a 598 amino acino acid precursor including a 21 aa signal sequence, a 440 aa extracellular domain (ECD), a 21 aa transmenbrane segment, and a 116 aa cytoplasmic domain. LIR-2 binds to

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Human Leukocyte Ig-Like Receptor B2/LILRB2/ILT4/CD85d (C-Avi-6His) Biotinylated



Catalog Number:PKSH033919

Classical MHCI proteins. Ligation of LIR-2 includes Tyr phosphorylation within its cytoplasmic ITIMs, a requirement for association with SHP-1. LIR-2 mediates tolerogenic DC-induced CD4+ T cell energy in vitro and in vivo.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com