

# Recombinant Human Placenta Growth Factor/PGF/PIGF/PLGF (C-6His)

Catalog Number:PKSH033937

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	PIGF2;PIGF-2;PGF;PLGF;PIGF2;PIGF;PGFL
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Leu19-Arg170
<b>Accession</b>	P49763-3
<b>Calculated Molecular Weight</b>	18.2 kDa
<b>Observed molecular weight</b>	25-30 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Placental growth factor is a protein that in humans is encoded by the PGF gene. It is a secreted protein and belongs to the PDGF/VEGF growth factor family. Alternate splicing results in at least three human mature PIGF forms containing 131 (PIGF?1), 152 (PIGF?2), and 203 (PIGF?3) amino acids (aa) respectively. PIGF is mainly found as a variably glycosylated, secreted, 55 ? 60 kDa disulfide linked homodimer. The protein is a member of the VEGF (vascular endothelial growth factor) sub-family-a key molecule in angiogenesis and vasculogenesis, in particular during embryogenesis. The main source of PGF during pregnancy is the placental trophoblast. PGF is also expressed in many other tissues, including the villous trophoblast. PIGF (especially PIGF?1) and some forms of VEGF can form dimers that decrease the angiogenic effect of VEGF on VEGF R2. PIGF?2, like VEGF164/165, shows heparin?dependent binding of

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human Placenta Growth Factor/PGF/PIGF/PLGF (C-6His)



Catalog Number:PKSH033937

---

neuropilin (Npn)<sup>1</sup> and Npn<sup>2</sup>, and can inhibit nerve growth cone collapse. Circulating PIGF often correlates with tumor stage and aggressiveness, and therapeutic PIGF<sup>2</sup> antibodies are being investigated for their ability to inhibit tumor growth and angiogenesis.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)