

# Recombinant Human Interleukin-2/IL-2 (C-6His-Avi) Biotinylated



Catalog Number:PKSH033973

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |   |
|------------------------------------|---|
| <b>Synonyms</b>                    | Interleukin-2;IL-2;T-Cell Growth Factor;TCGF;Aldesleukin;IL2  |
| <b>Species</b>                     | Human   |
| <b>Expression Host</b>             | HEK293 Cells  |
| <b>Sequence</b>                    | Ala21-Thr153  |
| <b>Accession</b>                   | P60568  |
| <b>Calculated Molecular Weight</b> | 18.3 kDa  |
| <b>Observed molecular weight</b>   | 16-18 kDa   |
| <b>Tag</b>                         | C-His-Avi   |
| <b>Bioactivity</b>                 | Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED <sub>50</sub> for this effect is 0.2 ng/ml. |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 90 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.             |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



> 90 % as determined by reducing SDS-PAGE.

## Background

Interleukin-2(IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system,belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017