Recombinant Human B7-2/CD86 (C-Avi-6His) Biotinylated

Catalog Number: PKSH033999



Note: Centrifuge before opening to ensure complete recovery of vial contents.

T	4.0
Descri	ntion
DCSCII	DOTOIL

Synonyms T-Lymphocyte Activation Antigen CD86;Activation B7-2

Antigen;B70;BU63;CTLA-4 Counter-Receptor B7.2;FUN-1;CD86;CD28LG2

Species Human

Expression Host HEK293 Cells
Sequence Ala24-Pro247
Accession AAH40261.1
Calculated Molecular Weight 28.4 kDa
Observed molecular weight 55-80 kDa
Tag C-Avi-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The protein is the receptor that involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation. The protein interacts with MARCH8, human herpesvirus 8 MIR2 protein, adenovirus subgroup B fiber proteins and acts as a receptor for these viruses.It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation.It contains 1 Ig-like C2-type(immunoglobulin-like)

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Human B7-2/CD86 (C-Avi-6His) Biotinylated 🍮

Catalog Number:PKSH033999



domainand 1 Ig-like V-type (immunoglobulin-like) domain.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>