

Recombinant Human B7-2/CD86 (C-Avi-6His) Biotinylated

Catalog No. PKSH033999

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	T-Lymphocyte Activation Antigen CD86;Activation B7-2 Antigen;B70;BU63;CTLA-4 Counter-Receptor B7.2;FUN-1;CD86;CD28LG2
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala24-Pro247
Accession	AAH40261.1
Calculated Molecular Weight	28.4 kDa
Observed molecular weight	55-80 kDa
Tag	C-Avi-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The protein is the receptor that involved in the costimulatory signal essential for T-lymphocyte proliferation and

For Research Use Only

interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation. The protein interacts with MARCH8, human herpesvirus 8 MIR2 protein, adenovirus subgroup B fiber proteins and acts as a receptor for these viruses. It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation. It contains 1 Ig-like C2-type (immunoglobulin-like) domain and 1 Ig-like V-type (immunoglobulin-like) domain.