

Recombinant Human CD3E&CD3G Heterodimer (C-Fc-6His&C-Fc-Flag)



Catalog Number:PKSH034015

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD3 epsilon & CD3 gamma;CD3E & CD3G;CD3E & CD3G Heterodimer;IMD18;T3E;TCRE
Species	Human
Expression Host	HEK293 Cells
Sequence	Asp23-Asp126&Gln23-Ser116
Accession	P07766&P09693
Calculated Molecular Weight	39.2&38.1 kDa
Observed molecular weight	40-60 kDa
Tag	C-Fc-Flag & C-Fc-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

T-cell surface glycoprotein CD3 epsilon & CD3 gamma chain, also known as CD3E & CD3G, are single-pass type I membrane proteins. CD3D, together with CD3- epsilon(CD3E) , CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017