# Recombinant Human CD68/SR-D1 (C-Fc)

Catalog Number: PKSH034029



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms CD68 antigenmacrophage antigen CD68;CD68

molecule;CD68;gp110;Macrosialin;SCARD1;SRD1;SR-D1

Species Human

Expression Host HEK293 Cells
Sequence Asn22-Ile320
Accession P34810
Calculated Molecular Weight 58.5 kDa
Observed molecular weight 70-100 kDa
Tag C-Fc

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

CD68, also called Scavenger Receptor D1 (SR-D1) and LAMP-4, is a heavily glycosylated type I transmembrane glycoprotein that belongs to the LAMP family. CD68 is expressed in many tumor cell lines which could allow them to attach to selectins on vascular endothelium, facilitating their dissemination to secondary sites. CD68 plays a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. It is a commonly used marker for macrophages. CD68 is also a biomarker of rheumatoid arthritis and Hodgkin's lymphoma. CD68 influences the functions of cells through NF-kB/focal adhesion kinase pathway (7). In addition, CD68 on macrophages binds tightly to both S100A8 and S100A9 to enhance the cell immunity.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com