# Recombinant Human IL-5 (C-6His-Avi) Biotinylated

Catalog Number:PKSH034042



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms nterleukin-5;IL-5;B-cell differentiation factor I;Eosinophil differentiation factor;T-

cell replacing factor;TRF;IL5

Species Human

Expression Host

Sequence

Ile20-Ser134

Accession

P05113

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Ile20-Ser134

P05113

Calculated Molecular Weight

16 kDa

18-24 kDa

C-His-Avi

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

IL-5 is expressed in eosinophils, NK cells, TC2 CD8+ T cells, mast cells, CD45+ CD4+ T cells, gamma delta T cells and IL-1 beta activated endothelial cells. IL-5 acts as a growth and differentiation factor for both B cells and eosinophils. Relative to B cells, IL-5 appears to induce the differentiation of activated conventional B-2 cells into Ig-secreting cells. In addition, it induces the growth of B-1 progenitors as well as IgM production by B-1 cells.IL-5 appears to perform a number of functions on eosinophils. These include the down modulation of Mac-1,the upregulation of receptors for IgA and IgG,the stimulation of lipid mediator (leukotriene C4 and PAF) secretion and the induction of granule release.IL-5 also promotes the growth and differentiation of eosinophils.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>