A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse CD7/Leu-9 (C-6His)

Catalog No. PKSM041424

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms T-Cell Antigen CD7;GP40;T-Cell Leukemia Antigen;T-Cell Surface Antigen

Leu-9;TP41;CD7

Species Mouse

Expression Host HEK293 Cells **Sequence** Gln24--Pro150

AccessionP50283Calculated Molecular Weight15.1 kDaObserved molecular weight18-30 kDaTagC-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 0.01 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



>95~% as determined by reducing SDS-PAGE.

Background

T-Cell Antigen CD7 is a single-pass type I membrane protein that that belongs to the the immunoglobulin superfamily.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Human CD7 is synthesized as a 240 amino acid precursor that contains a 25 amino acid signal sequence and a 215 amino acid mature chain with a Ig-like (immunoglobulin-like) domain. CD7 is normally expressed on all T-lymphocytes, NKcells, pre-B lymphocytes and pleuripotent hematopoietic stem cells. CD7 plays an essential role in T-cell interactions, Tcell/B-cell interaction during early lymphoid development, T- and NK-cell activation and cytokine production. CD7 has been shown to interact with PIK3R1and SECTM1. However, the function of the CD7 protein in the immune system is still largely unknown.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com