

PDHA1 Polyclonal Antibody

Catalog No. E-AB-60493

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse, Rat
Immunogen	Recombinant fusion protein of human PDHA1 (NP_000275.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Applications Recommended Dilution

WB	1:500-1:2000
IHC	1:50-1:200
IF	1:50-1:200


Data




Western blot analysis of extracts of various cell lines using PDHA1 Polyclonal Antibody at dilution of 1:1000.

Observed Mw:45kDa


Calculated Mw:40kDa/43kDa/44kDa/47kDa



Immunohistochemistry of paraffin-embedded Human colon carcinoma using PDHA1 Polyclonal Antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded Human mammary cancer using PDHA1 Polyclonal Antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded Human stomach using PDHA1 Polyclonal Antibody at dilution of 1:200 (40x lens).

For Research Use Only



Immunofluorescence analysis of NIH-3T3 cells using PDHA1 Polyclonal Antibody at dilution of 1:100.
Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using PDHA1 Polyclonal Antibody at dilution of 1:100.
Blue: DAPI for nuclear staining.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂, and provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle. The PDH complex is composed of multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase (E3). The E1 enzyme is a heterotetramer of two alpha and two beta subunits. This gene encodes the E1 alpha 1 subunit containing the E1 active site, and plays a key role in the function of the PDH complex. Mutations in this gene are associated with pyruvate dehydrogenase E1-alpha deficiency and X-linked Leigh syndrome. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.