

COX6A1 Polyclonal Antibody

Catalog No. E-AB-62050

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse, Rat
Immunogen	Recombinant fusion protein of human COX6A1 (NP_004364.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Applications Recommended Dilution


WB	1:200-1:2000
IHC	1:20-1:200
IF	1:50-1:200

Data




Western blot analysis of extracts of various cell lines using COX6A1 Polyclonal Antibody at dilution of 1:1000.


Observed Mw:12kDa
Calculated Mw:12kDa



Immunohistochemistry of paraffin-embedded Rat cerebellum using COX6A1 Polyclonal Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded Human liver cancer using COX6A1 Polyclonal Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded Mouse liver using COX6A1 Polyclonal Antibody at dilution of 1:100 (40x lens).

For Research Use Only



Immunofluorescence analysis of C6 cells using COX6A1 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using COX6A1 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Cytochrome c oxidase (COX), the terminal enzyme of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in the electron transfer and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 1 (liver isoform) of subunit VIa, and polypeptide 1 is found in all non-muscle tissues. Polypeptide 2 (heart/muscle isoform) of subunit VIa is encoded by a different gene, and is present only in striated muscles. These two polypeptides share 66% amino acid sequence identity. It has been reported that there may be several pseudogenes on chromosomes 1, 6, 7q21, 7q31-32 and 12. However, only one pseudogene (COX6A1P) on chromosome 1p31.1 has been documented.

For Research Use Only