

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse
Immunogen	Recombinant fusion protein of human AZGP1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Applications Recommended Dilution

WB	1:500-1:2000
-----------	--------------

Data



Western blot analysis of extracts of human serum using Zinc- α 2-glycoprotein (ZAG/Zinc- α 2-glycoprotein (ZAG/AZGP1)) Polyclonal Antibody at 1:1000 dilution.

Observed Mw:41kDa
Calculated Mw:34kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Zinc- α 2-Glycoprotein (AZGP1) can be found in blood plasma, seminal plasma, urine, sweat, saliva, liver, and epithelial cells of various human glands. AZGP1 has been proposed in the regulation of body weight, and the melanin production by normal and malignant melanocytes. AZGP1 stimulates lipid degradation in adipocytes and causes the extensive fat losses associated with some advanced cancers. AZGP1 has been reported to stimulate lipid breakdown and may have an important role in lipid homeostasis. Mature human AZGP1 consists of one MHC class I antigen region and a C2-type Ig-like domain. AZGP1 has two alternate splice forms, one shows a 66 amino acids substitution for the C-terminal 30 amino acids, the other one shows a nine Lys substitution for amino acid 151-298.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017