KAT7 Polyclonal Antibody

Catalog Number: E-AB-61134



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Recombinant fusion protein of human KAT7

Host Rabbit
Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

Formulation PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Applications Recommended Dilution


WB 1:500-1:2000 IHC 1:50-1:200

Data




Western blot analysis of extracts of various cell lines using KAT7 Polyclonal Antibody at 1:1000 dilution.


Observed Mw:63kDa Calculated Mw:51kDa/55kDa/58kDa/66kDa/70kDa



Immunohistochemistry of paraffin-embedded rat kidney using KAT7 Polyclonal antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded rat spleen using KAT7 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human colon carcinoma using KAT7 Polyclonal antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

For Research Use Only


Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>


KAT7 Polyclonal Antibody

Catalog Number: E-AB-61134





Immunohistochemistry of paraffin-embedded human breast cancer using KAT7 Polyclonal antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded mouse liver using KAT7 Polyclonal antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval wi

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene is part of the multimeric HBO1 complex, which possesses histone H4-specific acetyltransferase activity. This activity is required for functional replication origins and is involved in transcriptional activation of some genes. In both cases, the acetylation of histone H4 helps unfold chromatin so that the DNA can be accessed and replicated or transcribed.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com

Email: techsupport@elabscience.com