# **Elabscience**®

# PE/Cyanine7 Anti-Human CD83 Antibody[HB15e]

Catalog No.E-AB-F0993HStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityHumanApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

# **Antigen Information**

Alternate Names	BL11,CD83,HB15
Uniprot ID	Q01151
Gene ID	9308
Background	CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the
	immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells,
	and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen
	presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83
	is considered to be a useful marker for mature dendritic cells.

#### **Product Details**

Form	Liquid
Size	20Tests/100Tests/100Tests×2
Clone No.	HB15e
Host	Mouse
Isotype	Mouse IgG1, ĸ
Reactivity	Human
Application	FCM
Isotype Control	PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution.
	Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.
	This product is guaranteed up to one year from purchase.

**For Research Use Only** 

# **Elabscience**®

# Fluorophore


#### Conjugation: PE/Cyanine7

PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).

### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Product data**



PMA and ionomycin-stimulated (4h) Jurkat cells are stained with PE/Cyanine7 Anti-Human CD83 Antibody[HB15e] (filled gray histogram) or PE/Cyanine7 Mouse IgG1, κ Isotype Control (empty black histogram).

## **Related Information**

- 1. Sample Preparation for Flow Cytometry <u>https://www.elabscience.com/List-detail-5594.html</u>
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>