

MORF4L1 Polyclonal Antibody

Catalog No. E-AB-61505

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Recombinant fusion protein of human MORF4L1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Applications Recommended Dilution

WB 1:500-1:2000

Data



Western blot analysis of extracts of various cell lines using MORF4L1 Polyclonal Antibody at 1:1000 dilution.

Observed Mw:41kDa

Calculated Mw:26kDa/37kDa/41kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Component of the NuA4 histone acetyltransferase (HAT complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome-DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress transcription by deacetylation of nucleosomal histones. Required for homologous recombination repair (HRR and resistance to mitomycin C (MMC. Involved in the localization of PALB2, BRCA2 and RAD51, but not BRCA1, to DNA-damage foci.

For Research Use Only