

HMGB1 Polyclonal Antibody

Catalog No. E-AB-70044

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	KLH conjugated Synthetic peptide corresponding to Mouse HMGB1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 1% protective protein and 50% glycerol, pH7.4

Applications Recommended Dilution


WB	1:500-1:2000
IHC	1:200-1:800
IF	1:200-1:800

Data




Western Blot analysis of various samples using HMGB1 Polyclonal Antibody at dilution of 1:1000.


Observed Mw:25kDa
Calculated Mw:25kDa



Immunohistochemistry analysis of paraffin-embedded Human liver cancer using HMGB1 Polyclonal Antibody at dilution of 1:500.




Immunohistochemistry analysis of paraffin-embedded mouse kidney using HMGB1 Polyclonal Antibody at dilution of 1:500.



Immunofluorescence analysis of paraffin-embedded Mouse heart using HMGB1 Polyclonal Antibody at dilution of 1:300.

For Research Use Only



Immunohistochemistry analysis of paraffin-embedded Human Tonsil using HMGB1 Polyclonal Antibody(Elabscience® Product Detected by Lifespan).

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

High mobility group (HMG) proteins 1 and 2 are ubiquitous non-histone components of chromatin. Evidence suggests that the binding of HMG proteins to DNA induces alterations in the DNA architecture including DNA bending and unwinding of the helix. HMG proteins synergize with Oct-2, members of the NF B family, ATF-2 and c-Jun to activate transcription. Other studies indicate that phosphorylation of HMG protein is required to stimulate the transcriptional activity of the protein. Human HMG-1 and HMG-2 both contain two DNA-binding domains, termed HMG boxes. HMG proteins bind single-stranded DNA but induce conformational changes in double-stranded DNA alone.