

BrdU Monoclonal Antibody

Catalog No. E-AB-70248

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	KLH conjugated BrdU
Host	Mouse
Isotype	IgG
Clone	4B6C12
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 1% protective protein and 50% glycerol, pH7.4


Applications Recommended Dilution

IHC	1:100-1:500
IF	1:100-1:500

Data



Immunohistochemistry analysis of paraffin-embedded mouse jejunum (intraperitoneal injection BrdU every 2 hours for 4 times) using BrdU Monoclonal Antibody at dilution of 1:300.



Immunofluorescence analysis of HeLa cells (treated with 0.03mg/ml BrdU for 40 min) using BrdU Monoclonal Antibody at dilution of 1:100.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Bromodeoxyuridine (5-bromo-2'-deoxyuridine, BrdU, BUdR, BrdUrd) is a synthetic nucleoside that is an analog of thymidine. It can be incorporated into the newly synthesized DNA of replicating cells (during the S phase of the cell cycle), substituting for thymidine during DNA replication. As such, BrdU is used for birth dating and monitoring cell proliferation. BrdU is a toxic and mutagenic substance. It triggers cell death, the formation of teratomas, alters DNA stability, lengthens the cell cycle, and has mitogenic, transcriptional and translational effects on cells that incorporate it.

For Research Use Only