Recombinant SARS-CoV-2 S-stable trimer Protein (C-6His)

Catalog Number:PKSV030314

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	SARS-CoV-2 Spike;SARS-CoV-2 Spike glycoprotein;SARS-CoV-2 S glycoprotein;SARS-CoV-2 S-stable trimer Protein
Species	SARS-CoV-2
Expression Host	HEK293 Cells
Accession	QHD43416.1
Calculated Molecular Weight	136.6 kDa
Observed molecular weight	170-220 kDa
Tag	C-His
Bioactivity	Immobilized 2019-nCoV S-trimer Protein-His(PKSV030314) at 2μg/ml (100 μl/well) can bind Human ACE-2-Fc(PKSR030492). The ED50 of Human ACE-2-Fc(PKSR030492) is 23.83 ng/ml, Immobilized 2019-nCoV S-trimer Protein-His(PKSV030314) at 2μg/ml (100 μl/well) can bind Anti-2019-nCoV S1 mAb (5D9). The ED50 of Anti-2019-nCoV S1 mAb (5D9) is 15.44 ng/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 μm filtered solution of PBS, 5% Trehalose, 10% Glycerol, 5% Mannitol, 0.01% Tween 80, pH 7.4.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com