

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

PE/Cyanine7 Anti-Mouse CD73 Antibody[TY/23]

Catalog No.E-AB-F1089UHReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names 5'-nucleotidase,Nt5e,5'-NT,Ecto-5'-nucleotidase,CD73,Nt5, Nte,Nt5e

Uniprot ID Q61503 Gene ID 23959

Background CD73 (ecto-5`-nucleotidase) is a 69 kD GPI-anchored surface protein. In mice, expression of

CD73 in bone marrow is restricted to CD11b+ myeloid cells. In spleen, it is largely expressed on

T cells.

Product Details

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.2 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & TY/23 \\ Host & Rat \\ \end{tabular}$

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG2a, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & FCM \\ \end{tabular}$

Isotype Control PE/Cyanine 7 Rat IgG2a, K Isotype Control [2A3] [Product E-AB-F09833H]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Tor Research Ose Only

Fax: 1-832-243-6017

A Reliable Research Partner in Life Science and Medicine

Fluorophore


Conjugation: PE/Cyanine7

PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in $100~\mu L$ volume].

Product data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD3 Antibody and PE/Cyanine7 Anti-Mouse CD73 Antibody (Left). Splenocytes are stained with FITC Anti-Mouse CD3 Antibody and PE/Cyanine7 Rat IgG2a, κ Isotype Control (Right).

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com