



A Reliable Research Partner in Life Science and Medicine

PE/Elab Fluor® 594 Anti-Mouse CD19 Antibody[1D3]

Catalog No.E-AB-F0986UPReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names B-lymphocyte antigen CD19,Cd19,Differentiation antigen CD19,CD19

Uniprot ID P25918 **Gene ID** 12478

Background CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed

on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex

integral to B cell activation.

Product Details

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.2 \ mg/mL \\ Size & 25 \mu g/100 \mu g \\ \end{tabular}$

Clone No. 1D3 Host Rat

IsotypeRat IgG2a, κReactivityMouseApplicationFCM

Isotype Control PE/Elab Fluor[®] 594 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833P]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

A Reliable Research Partner in Life Science and Medicine

Fluorophore


Conjugation: PE/Elab Fluor® 594

PE/Elab Fluor[®] 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in $100~\mu L$ volume].

Product data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] Red 780 Anti-Mouse CD45R/B220 Antibody and PE/Elab Fluor[®] 594 Anti-Mouse CD19 Antibody (Left). Splenocytes are stained with Elab Fluor[®] Red 780 Anti-Mouse CD45R/B220 Antibody and PE/Elab Fluor[®] 594 Rat IgG2a, κ Isotype Control (Right).

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com