

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 488 Anti-Mouse Perforin Antibody[S16009A]

Catalog No.E-AB-F1294LReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsICFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

Alternate Names Prf1, Pfp, Pfn, Pfn-1, Cytolysin, Lymphocyte pore-forming protein

Uniprot ID P10820 Gene ID 18646

**Background** Perforin is a 70 kD cytolytic protein that is expressed in the cytoplasmic granules of cytotoxic T

lymphocytes (CTLs) and natural killer (NK) cells. Perforin is one of the major effector molecules

used by cytotoxic T cells and NK cells to mediate targeted cell lysis.

#### **Product Details**

Form Liquid

Size 50Tests/100Tests/100Tests×2

Clone No. S16009A Host Rat

IsotypeRat IgG2a, κReactivityMouseApplicationICFCM

**Isotype Control** Elab Fluor<sup>®</sup> 488 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832L]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

#### **Elabscience Bionovation Inc.**

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# **Fluorophore**

Conjugation: Elab Fluor® 488

Elab Fluor<sup>®</sup> 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

### Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086