



A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 488 Anti-Mouse Perforin Antibody[S16009A]

Catalog No.E-AB-F1294ULReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsICFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Prf1, Pfp, Pfn, Pfn-1, Cytolysin, Lymphocyte pore-forming protein

Uniprot ID P10820 **Gene ID** 18646

Background Perforin is a 70 kD cytolytic protein that is expressed in the cytoplasmic granules of cytotoxic T

lymphocytes (CTLs) and natural killer (NK) cells. Perforin is one of the major effector molecules

used by cytotoxic T cells and NK cells to mediate targeted cell lysis.

Product Details

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.5 mg/ml \\ Size & 25 \mu g/100 \mu g \\ Clone No. & S16009A \\ Host & Rat \\ \end{tabular}$

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG2a, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & ICFCM \\ \end{tabular}$

Isotype Control Elab Fluor® 488 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833L]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Elabscience Bionovation Inc.

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: Elab Fluor® 488

Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶ cells in 100 μL volume].

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086