



A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Mouse TCRβ Antibody[H57-597 (HB218)]

Catalog No.E-AB-F1123QReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names TCR-β chain, TCR-β, β-TCR

Gene ID 21577

Background T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ

chain (TCR γ/δ). TCR- β is a member of the immunoglobulin superfamily and a component of the

CD3/TCR complex (along with TCR- α). It is expressed on α/β TCR-bearing T cells and

thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction,

and T cell activation.

Product Details

Form Liquid

Size 50Tests/100Tests/100Tests×2

Clone No. H57-597 (HB218)

Host Armenian Hamster

Isotype Armenian Hamster IgG

Reactivity Mouse **Application** FCM

Isotype Control Storage BufferElab Fluor® Violet 450 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852Q]
Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Fax: 1-832-243-6017

A Reliable Research Partner in Life Science and Medicine

Fluorophore


Conjugation: Elab Fluor® Violet 450

Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Product data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] Violet 450 Anti-Mouse TCRβ Antibody (filled gray histogram) or Elab Fluor[®] Violet 450 Armenian hamster IgG Isotype Control (empty black histogram).

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com