# **Elabscience**®

# Elab Fluor<sup>®</sup> Violet 450 Anti-Mouse TCRβ Antibody[H57-597 (HB218)]

Catalog No.E-AB-F1123UQStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityMouseApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

#### **Antigen Information**

Alternate Names	TCR- $\beta$ chain, TCR- $\beta$ , $\beta$ -TCR
Gene ID	21577
Background	T cell receptor (TCR) is a heterodimer consisting of an $\alpha$ and a $\beta$ chain (TCR $\alpha/\beta$ ) or a $\gamma$ and a $\delta$
	chain (TCR $\gamma/\delta$ ). TCR- $\beta$ is a member of the immunoglobulin superfamily and a component of the
	CD3/TCR complex (along with TCR- $\alpha$ ). It is expressed on $\alpha/\beta$ TCR-bearing T cells and
	thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction,
	and T cell activation.

#### **Product Details**

Form	Liquid
Concentration	0.5 mg/mL
Size	25µg/100µg
Clone No.	H57-597 (HB218)
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Reactivity	Mouse
Application	FCM
Isotype Control	Elab Fluor <sup>®</sup> Violet 450 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853Q]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution.
	Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.
	This product is guaranteed up to one year from purchase.

**For Research Use Only** 

# **Elabscience**®

# Fluorophore


Conjugation: Elab Fluor<sup>®</sup> Violet 450

Elab Fluor<sup>®</sup> Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is  $0.1-1 \ \mu g/10^6$  cells in  $100 \ \mu L$  volume].

## **Product data**



C57BL/6 murine splenocytes are stained with Elab Fluor<sup>®</sup> Violet 450 Anti-Mouse TCRβ Antibody (filled gray histogram) or Elab Fluor<sup>®</sup> Violet 450 Armenian hamster IgG Isotype Control (empty black histogram).

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <u>https://www.elabscience.com/List-detail-5594.html</u>
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>