



A Reliable Research Partner in Life Science and Medicine

## **Recombinant Human FGF-18 protein(His Tag)**

Catalog No. PKSH034159

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms FGFI;zFGF5
Species Human
Expression Host E.coli

Sequence Ala 27-Arg 199

AccessionO76093Calculated Molecular Weight21.1 kDaObserved molecular weight20 kDaTagC-His

**Bioactivity** Measure by its ability to induce 3T3 cells proliferation. The  $ED_{50}$  for this effect is

1.3-2.0 ng/mL. The specific activity of recombinant human FGF-18 is  $> 5 \times 10^5$ 

IU/mg.

## **Properties**

**Purity** > 98 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 0.1 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS,pH 8.0.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## **Background**

Fibroblast growth factor 18 (FGF18) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. It has been shown in vitro that FGF18 is able to induce neurite outgrowth in PC12 cells. Studies of the similar proteins in mouse and chick suggested that this protein is a pleiotropic growth factor that stimulates proliferation in a number of tissues, most notably the liver and small intestine. Experiment datas identified FGF18 as a selective ligand for FGFR3 in limb bud mesenchymal cells, which suppressed proliferation and promoted their differentiation and production of cartilage matrix. FGF18 appears to regulate cell proliferation and differentiation positively in osteogenesis and negatively in chondrogenesis.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u>

Email: techsupport@elabscience.com