A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse IL-7 protein(His Tag)

Catalog No. PKSM041462

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Interleukin-7;IL-7;IL7

SpeciesMouseExpression HostE.coli

Sequence Glu 26-Ile 154

Accession P10168
Calculated Molecular Weight 15.8 kDa
Observed molecular weight 16 kDa
Tag C-His

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Mouse interleukin-7(IL-7) is the member of hemopoietin family which is important to the differentiation, proliferation, and survival of lymphocyte. Mouse IL-7 shares approximately 88% as sequence identity with rat IL-7 and 58-60% with human, equine, bovine, ovine, porcine, feline and canine IL-7. It is widely expressed in primary and secondary lymphoid

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

tissues cell and stromal epithelial cells of the thymus, bone marrow, and intestines. IL-7 activation of IL-7 R alpha is critical for both T cell and B cell lineage development. It is important for proliferation during certain stages of B-cell maturation. IL-7 contributes to the maintenance of all na?ve and memory T cells, mainly by promoting expression of the anti-apoptotic protein Bcl-2. It is required for optimal T cell-dendritic cell interaction.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com