
## **Recombinant Mouse IL-17B protein(His Tag)**

Catalog Number: PKSM041494



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                 |   |
|-----------------------------|---|
| Synonyms                    | TNFSF1B;Lymphotoxin-alpha (LT-α);LTalpha;Ltx;TNF-<br>be;TNFSF1;Tnf;Tnfb;Tnfsf;Tnfsf1b;Tnlg1e;hlb38;hlb382;lymph   |
| Species                     | Mouse   |
| Expression Host             | E.coli  |
| Sequence                    | His 21-Phe 180  |
| Accession                   | Q9QXT6  |
| Calculated Molecular Weight | 19.0 kDa  |
| Observed molecular weight   | 17-25 kDa   |
| Tag                         | C-His   |
| Bioactivity                 | Measure by its ability to induce IL-8 secretion in HepG2 cells. The $ED_{50}$ for this effect is < 1.5 ng/mL.   |
| Properties                  |   |
| Purity                      | > 98 % as determined by reducing SDS-PAGE.  |
| Endotoxin                   | < 0.1 EU per µg of the protein as determined by the LAL method.   |
| Storage                     | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping                    | This product is provided as lyophilized powder which is shipped with ice packs.   |
| Formulation                 | Lyophilized from sterile PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as<br>protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.        |
| Reconstitution              | Please refer to the printed manual for detailed information.  |
| Data                        |   |



> 98 % as determined by reducing SDS-PAGE.

## Background

Interleukin-17B (IL-17B) is a member of IL-17 cytokines family that plays important roles in host defence responses and inflammatory diseases. In addition to IL-17B, members of the IL-17 family include IL-17A, IL-17C, IL-17D, IL-17E and IL-17F. The six IL-17 cytokines are highly conserved at C terminus, and contain five spatially conserved cysteine residues that mediate dimerization. Expression of IL-17B protein has been reported in neurons, testis, ovary, stomach, pancreas, small intestine and chondrocytes. IL-17B binds to Interleukine-17 receptor B (IL-17 RB) and induces the production of

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Catalog Number:PKSM041494



inflammatory cytokines and the infiltration of inflammatory immune cells.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com