

Recombinant Swine IFN gamma protein(His Tag)

Catalog Number:PKSS000012



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|---|
| Synonyms | IFN-gamma;IFNG |
| Species | Porcine |
| Expression Host | E.coli |
| Sequence | Gln 24-Lys 166 |
| Accession | P17803 |
| Calculated Molecular Weight | 17.7 kDa |
| Observed molecular weight | 15 kDa |
| Tag | C-His |
| Bioactivity | Measure by its ability to protect PK15 cells infected with encephalomyocarditis (EMC) virus.The ED ₅₀ for this effect is < 40 pg/mL. |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.01 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

IFN γ is the major interferon produced by mitogenically or antigenically stimulated lymphocytes. It is structurally different from type I interferon and its major activity is immunoregulation. It has been implicated in the expression of class II histocompatibility antigens in cells that do not normally produce them; leading to autoimmune disease. Interferon gamma is produced mainly by T-cells and natural killer cells activated by antigens; mitogens; or alloantigens. It is produced by lymphocytes expressing the surface antigens CD4 and CD8. IFN γ synthesis is induced by IL-2; FGF-basic; and EGF.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017