



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD5 Antibody[53-7.3]

Catalog No. E-AB-F1185B Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM**

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Ly-1, Lyt-1, Lymphocyte antigen 1, CD5, Cd5

Uniprot ID P13379 Gene ID 12507

Background CD5 is a 67 kD protein, also known as Lyt-1, Ly-1, T1, Tp67, or Ly-12. It is a member of the

> scavenger receptor cysteine-rich protein superfamily (SRCR) and primarily expressed on thymocytes, T cells, and B-1 cells. Although mature α/β T cells express high levels of CD5, very few γ/δ T cells express this antigen. The interaction of CD5 with CD72, gp35-37, TCR, or BCR

is involved in T and B cell activation.

Product Details

Form Liquid Concentration 0.5 mg/mL Size $25 \mu g / 100 \mu g$ Clone No. 53-7.3 Host Rat

Rat IgG2a, κ **Isotype** Reactivity Mouse **Application FCM**

Isotype Control Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution. Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Email: techsupport@elabscience.com Web: www.elabscience.com


Tel: 1-832-243-6086 Fax: 1-832-243-6017

A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0~\mu g$ per 10^6 cells in $100~\mu L$ volume or $100~\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Product data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD45R/B220 Antibody and Biotin Anti-Mouse CD5 Antibody followed by Streptavidin-Elab Fluor[®] 647 (Left). Splenocytes are stained with FITC Anti-Mouse CD45R/B220 Antibody and Biotin Rat IgG2a, κ Isotype Control followed by Streptavidin-Elab Fluor[®] 647 (Right).

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com