

# Recombinant Human B7-H4/VTCN1 Protein (aa 29-258, His Tag)



Catalog Number: PDMH100005

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	B7S1;B7x;Vtcn1;B7h.5;B7-H4;B7H4T-cell costimulatory molecule B7x;B7S1VCTN1;B7XPRO1291;FLJ22418;Immune costimulatory protein B7-H4;Protein B7S1;T cell costimulatory molecule B7x;V-set domain containing T cell activation inhibitor 1;V-set domain-containing T-cell activation inhibitor 1
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Phe29-Ala258
<b>Accession</b>	Q7Z7D3
<b>Calculated Molecular Weight</b>	26.8 kDa
<b>Observed molecular weight</b>	43-48 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

V-set domain-containing T-cell activation inhibitor 1; also known as B7X; B7H4; B7S1; and VTCN1; is a single-pass type? membrane protein belonging to the B7 family of costimulatory proteins. These proteins are expressed on the surface of antigen-presenting cells and interact with ligands on T lymphocytes. They provide costimulatory signals that regulate T cell responses. A soluble form of B7H4 has also been detected. B7X / VTCN1 / B7H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation; proliferation; cytokine production and development of

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human B7-H4/VTCN1 Protein (aa 29-258, His Tag)



Catalog Number: PDMH100005

---

cytotoxicity. When expressed on the cell surface of tumor macrophages; B7X / VTCN1 / B7H4 plays an important role; together with regulatory T-cells(Treg); in the suppression of tumor-associated antigen-specific T-cell immunity. B7X / VTCN1 / B7H4 is also involved in promoting epithelial cell transformation. This membrane protein can be up-regulated by IL6 / interleukin-6 and IL10 / interleukin-10 and inhibited by CSF2 / GM-CSF and IL4 / interleukin-4 on antigen-presenting cells.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)