



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD16 Antibody[3G8]

Catalog No. E-AB-F1236A Reactivity Human Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications** FCM.Block

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names FcγRIII,Fc gamma receptor,Fc gamma receptor 3

Uniprot ID P08637,O75015

Gene ID 941

Background CD16 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also

known as B7-1, B7, and Ly-53. CD16 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD16 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD16 appears to be expressed later in the immune response than CD86. CD16 can also bind to CD152, also known as CTLA-4, to deliver an

inhibitory signal to T cells.

Product Details

Form Liquid Concentration 0.5 mg/mL Size $25\mu g/100\mu g$

Clone No. 3G8 Host Mouse

Isotype Mouse IgG1, κ Reactivity Human **Application** FCM,Block

Isotype Control Purified Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793A]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5~\mu g$ per 10^6 cells in $100~\mu L$ volume or $100~\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017