# **Elabscience**®

## APC Anti-Rat CD45RA Antibody[OX-33]

Catalog No.E-AB-F1306EStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityRatApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

Alternate Names	Leukocyte common antigen, high molecular weight form
Gene ID	24699
Background	CD45 is a protein tyrosine phosphatase with multiple isoforms differing as a result of alternative
	splicing of the extracellular domain and glycosylation. CD45 is expressed on all hematopoietic
	cells except erythrocytes and platelets. CD45RA is one of the CD45 isoforms with a molecular
	weight of 200-220 kD. It is expressed almost exclusively on B cells. CD45 functions in signal
	transduction through T and B cell antigen receptors. CD45 has been shown to interact with
	various proteins, including galectin-1, CD2, CD3, and CD4.

### **Product Details**

Form	Liquid
Size	50Tests/100Tests/100Tests×2
Clone No.	OX-33
Host	Mouse
Isotype	Mouse IgG1, κ
Reactivity	Rat
Application	FCM
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution.
	Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.
	This product is guaranteed up to one year from purchase.

**For Research Use Only** 

## **Elabscience**®

## Fluorophore


#### **Conjugation:** APC

APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Product data**



Rat splenocytes are stained with APC Anti-Rat CD45RA Antibody (filled gray histogram) or APC Mouse IgG1,  $\kappa$ Isotype Control (empty black histogram).

## **Related Information**

- 1. Sample Preparation for Flow Cytometry <u>https://www.elabscience.com/List-detail-5594.html</u>
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>