



A Reliable Research Partner in Life Science and Medicine

# Purified Anti-Human CD5 Antibody[HISM2]

Catalog No.E-AB-F1313AReactivityHumanStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

Alternate Names T-cell surface glycoprotein CD5,CD5,Lymphocyte antigen T1/Leu-1,LEU1

**Uniprot ID** P06127 **Gene ID** 921

**Background** CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member

of the scavenger receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions upon binding to ligands such

as CD72.

#### **Product Details**

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.5 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & HISM2 \\ Host & Mouse \\ Isotype & Mouse IgG1, <math>\kappa$  \\ \end{tabular}

**Reactivity** Human **Application** FCM

**Isotype Control** Purified Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793A]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping
Biological ice pack at 4 °C
Stability & Storage
Keep as concentrated solution.
Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>

Fax: 1-832-243-6017

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \,\mu g$  per  $10^6$  cells in  $100 \,\mu L$  volume or  $100 \,\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017